arXiv:1208.3833v1 [hep-th] 19 Aug 2012

HU-EP-12/19

Quantization of AdS×S particle in static gauge

George Jorjadze,[a,b] Chrysostomos Kalousios,[a] Zurab Kepuladze,[c]

[a]*Institut für Physik der Humboldt-Universität zu Berlin,*
Newtonstraße 15, D-12489 Berlin, Germany

[b]*RMI & Free University of Tbilisi,*
Bedia Str., 0183, Tbilisi, Georgia

[c]*ITP, Ilia State University & Andronikashvili Institute of Physics,*
Tamarashvili Str. 6, 0177, Tbilisi, Georgia

Abstract

We quantize the particle dynamics in $\mathrm{AdS}_{N+1}\times\mathrm{S}^M$ spacetime in static gauge, which leads to the coordinate representation with wave functions depending only on spatial coordinates. The energy square operator is quadratic in canonical momenta and contains a scalar curvature term. We analyze the self-adjointness of this operator and calculate its spectrum. We then construct unitary representations of the isometry group $\mathrm{SO}(2,N)\times\mathrm{SO}(M+1)$ and calculate the quantum relation between the Casimir numbers.

Introduction

In this paper we quantize the AdS×S particle dynamics in static gauge. We use the quantization scheme of [1] based on a covariant construction of the energy square operator in the coordinate representation, where the wave functions depend only on spatial coordinates. Let us first outline the scheme, which we present here in a slightly modified form.

Particle dynamics in a spacetime with coordinates x^μ, $\mu = (0, 1, ..., \mathcal{N})$ and a metric tensor $g_{\mu\nu}(x)$ can be described by the following action in the first order formalism

$$S = \int d\tau \Big(p_\mu \, \dot{x}^\mu - \frac{\lambda}{2} \left(g^{\mu\nu} p_\mu p_\nu + \mathcal{M}^2 \right) \Big) \,. \tag{1}$$

Here $\mathcal{M}$ is the particle mass, λ is a Lagrange multiplier and its variation provides the mass-shell condition

$$g^{\mu\nu} p_\mu p_\nu + \mathcal{M}^2 = 0 \,. \tag{2}$$

Using the Faddeev-Jackiw reduction [2] in the gauge

$$x^0 + p_0 \tau = 0 \,, \tag{3}$$

from (1) we get an ordinary Hamiltonian system[1]

$$S = \int d\tau \Big(p_n \, \dot{x}^n - \frac{1}{2} \, p_0^2 \Big) \,, \tag{4}$$

where p_0^2, as a function of the spatial coordinates and momenta (x^n, p_n) $(n = 1, ..., \mathcal{N})$, is obtained from the constraint (2) and the gauge fixing condition (3).

Notice that here we have modified the form of the standard static gauge $x^0 = \tau$ used in [1]. However, this modification does not change the quantization scheme and it appears more convenient for a Hamiltonian treatment of (4) in a static spacetime, as well as for the generalization to string theory [3].

A static spacetime metric tensor can be represented in the form

$$g_{\mu\nu} = \begin{pmatrix} g_{00} & 0 \\ 0 & g_{mn} \end{pmatrix} \,, \tag{5}$$

where g_{00} and g_{mn} are functions only of the spatial coordinates x^n. In this case the particle energy $E(p, x) = -p_0 > 0$ is conserved and from (2) follows that

$$E^2 = \Lambda(x) \, g^{mn}(x) \; p_m p_n + \mathcal{M}^2 \, \Lambda(x) \,, \tag{6}$$

with $\Lambda(x) := -g_{00}(x) > 0$.

Thus, the Hamiltonian in (4) corresponds to a motion of a particle in the potential field $V(x) = \frac{1}{2} \, \mathcal{M}^2 \, \Lambda(x)$ and in a curved background with metric tensor

$$h_{mn}(x) = \frac{1}{\Lambda(x)} \; g_{mn}(x) \,. \tag{7}$$

---

[1] We neglect the total derivative term $-\frac{\mathrm{d}}{\mathrm{d}\tau} \left(\frac{1}{2} \, p_0^2 \tau \right)$.

It is natural to quantize this system in the coordinate representation, where the wave functions $\psi(x)$ form a Hilbert space with covariant scalar product

$$\langle\psi_2|\psi_1\rangle = \int d^{\mathcal{N}}x\,\sqrt{h(x)}\;\psi_2^*(x)\,\psi_1(x)\;, \qquad\qquad h(x) := \det h_{mn}(x)\;. \tag{8}$$

On the basis of DeWitt's construction for quadratic in momenta operators [4], it was argued in [1] that the energy square operator is given by

$$E^2 = -\Delta_h + \frac{\mathcal{N}-1}{4\mathcal{N}}\,\mathcal{R}_h(x) + \mathcal{M}^2\,\Lambda(x)\;. \tag{9}$$

Here Δ_h is the covariant Laplace-Beltrami operator for the metric tensor h_{mn} and $\mathcal{R}_h(x)$ denotes the corresponding scalar curvature. The solution of the eigenvalue problem for (9) then provides the energy operator in diagonal form.

The coefficient in front of the scalar curvature term has been a subject of discussions during decades (see [5] and references therein). Therefore it is useful to comment on the value of this coefficient, $\frac{\mathcal{N}-1}{4\mathcal{N}}$, chosen in (9).

For the particle dynamics in AdS$_{\mathcal{N}+1}$ this coefficient was calculated in [1] from the commutation relations of the symmetry generators. In this case $\mathcal{R}_h(x)$ corresponds to the curvature of a semi-sphere and, therefore, it is constant. The obtained constant shift in the energy square operator provides its spectrum in the form $(E_0+n)^2$, with fixed E_0 and a non-negative integer n, that leads to the correct energy spectrum for the AdS particle.

For a generic $\mathcal{N}+1$ dimensional static spacetime the same value of the coefficient $\frac{\mathcal{N}-1}{4\mathcal{N}}$ follows from the equivalence between the static gauge quantization and the covariant quantization based on the Klein-Gordon type equation.

The covariant quantization is a more conventional approach to the AdS×S particle dynamics [6]. The general case with arbitrary dimensions and radii in this approach was analyzed in [7] from the perspective of scalar field propagators. A fixed gauge approach to the AdS$_5\times$ S^5 particle dynamics was considered in [8] in the context of string theory.

Our aim here is to apply the static gauge quantization scheme to AdS$_{N+1}\times$S^M particle. AdS$_{N+1}$ will be realized as a hyperbola $X^A\,X_A = -R^2,$ with $A = (0', 0, 1, ..., N)$, embedded in $\mathbb{R}^{2,\,N}$ and S^M as a M-dimensional sphere $Y_I\,Y_I = R_S^2$ in $\mathbb{R}^{M+1}$, with $I = (1, ..., M+1)$.

In the next section we deal with the classical case, preparing the system for quantization.

Geometry on AdS$_{N+1}\times$S^M

AdS$_{N+1}\times$S^M has $\mathcal{N} = N + M$ spatial coordinates and we associate the first N coordinates with the AdS part and the rest M coordinates with S^M. The time coordinate x^0 is given by the polar angle θ in the $(X^{0'}, X^0)$ plane.

We parameterize the embedding coordinates of $\mathbb{R}^{2,N}$ as follows

$$X^{0'} = \frac{R}{\rho}\,\sin\theta\;, \qquad X^0 = \frac{R}{\rho}\,\cos\theta\;, \qquad X^a = \frac{R}{\rho}\,x^a\;, \qquad \rho := \sqrt{1 - x^b x^b}\;, \tag{10}$$

where the coordinates x^a are given on the N-dimensional unit disk.

The choice of coordinates ϕ^α on S^M is not important for our calculations, since the quantization of the spherical part is trivial. We use the Greek letters (α, β) for tensorial indices on S^M and they run from $N+1$ to $N+M$.

The induced metric tensor on $\mathrm{AdS}_{N+1}\times\mathrm{S}^M$ has the structure (5) with

$$g_{00} = -\frac{R^2}{\rho^2}\,, \qquad g_{mn} = \begin{pmatrix} g_{ab} & 0 \\ 0 & g_{\alpha\beta} \end{pmatrix}\,, \tag{11}$$

where g_{ab} corresponds to the spatial part of AdS_{N+1}

$$g_{ab} = \frac{R^2}{\rho^2}\left(\delta_{ab} + \frac{x^a\,x^b}{\rho^2}\right) \tag{12}$$

and $g_{\alpha\beta}$ is the metric tensor on S^M. Note that the scalar curvature for g_{mn} is given by

$$\mathcal{R}_g = -N(N-1)\,R^{-2} + M(M-1)\,R_S^{-2}\,. \tag{13}$$

It is worth mentioning that the metric tensor (12) corresponds to the Euclidean AdS space (Lobachevski plane) and after the rescaling (7) with the Weyl factor

$$\frac{1}{\Lambda(x)} = \frac{\rho^2}{R^2}\,, \tag{14}$$

it becomes the metric tensor on the unit semi-sphere (see Fig. 1) with

$$h_{ab}(x) = \delta_{ab} + \frac{x^a\,x^b}{\rho^2}\,, \qquad \text{and} \qquad \det h_{ab} = \frac{1}{\rho^2}\,. \tag{15}$$

The total background metric tensor h_{mn} defined by (7) has a similar to (11) block structure and from (15) we find the integration measure in (8) to be

$$\sqrt{h} = \kappa^{-M}\,\mu_S\,\rho^{M-1}\,, \tag{16}$$

where $\kappa = R/R_S$ and μ_S is the $\mathrm{SO}(M+1)$ invariant measure on the unit sphere.

The Laplace-Beltrami operator for the metric tensor h_{mn} then reads

$$\Delta_h = h^{ab}\,\partial_{ab} - (N+M)x^a\,\partial_a + \frac{\kappa^2}{\rho^2}\,\Delta_S\,, \tag{17}$$

where

$$h^{ab} = \delta_{ab} - x^a\,x^b \tag{18}$$

is the inverse to h_{ab} and Δ_S is the Laplace-Beltrami operator on the unit sphere.

To calculate $\mathcal{R}_h$, one can use the transformation rule (A.1) of scalar curvatures under Weyl rescalings. With the help of (13) it leads to

$$\mathcal{R}_h = (N+M)\,(N+M-1) + \frac{M(M-1)\,(\kappa^2-1)}{\rho^2}\,. \tag{19}$$

Now we consider the dynamical integrals related to the $\mathrm{SO}(2,N)\times\mathrm{SO}(M+1)$ isometry transformations

$$J_{AB}=\mathcal{V}^a_{AB}\,p_a+\mathcal{V}^0_{AB}\,p_0\,, \qquad L_{IJ}=\mathcal{V}^\alpha_{IJ}\,\pi_\alpha\ . \tag{20}$$

Here p_a and π_α are the canonically conjugated variables to x^a and ϕ^α, respectively, p_0 is the negative square root of

$$E^2=\frac{R^2}{\rho^2}\left[g^{ab}(x)\,p_ap_b+g^{\alpha\beta}(\phi)\,\pi_\alpha\pi_\beta+\mathcal{M}^2\right] \tag{21}$$

and the coefficients of the momentum variables are the components of the Killing vector fields (see (A.3) in Appendix)

$$\mathcal{V}^0_{AB}=g^{00}(X_B\,\partial_\theta X_A-X_A\,\partial_\theta X_B)\ , \qquad \mathcal{V}^a_{AB}=g^{ab}(X_B\,\partial_b X_A-X_A\,\partial_b X_B)\ , \tag{22}$$

$$\mathcal{V}^\alpha_{IJ}=g^{\alpha\beta}(Y_J\,\partial_\beta Y_I-Y_I\,\partial_\beta Y_J)\ . \tag{23}$$

Since we use dimensionless coordinates, the momentum variables p_a, π_α and the energy are also dimensionless.[2] The minimal energy corresponds to the vanishing momenta and the maximal value of ρ (i.e. $\rho=1$), that yields $E^2_{min}=\mathcal{M}^2\,R^2$.

The vector field components for the boost generators in (22) depend on the time coordinate θ and to use them in (20), one has to make the replacement $\theta\mapsto E\tau$ corresponding to the gauge fixing (3). This calculation for the boost generators at $\tau=0$ yields

$$J_{a0'}=Ex^a\ , \qquad J_{a0}=(p_b\,x^b)x^a-p_a\ . \tag{24}$$

From the canonical Poisson brackets $\{p_a,\,x^b\}=\delta_{ab}$ follows that the energy square (21) and the boosts (24) satisfy the Poisson bracket relations

$$\{E^2,\,J_{a0'}\}=-2EJ_{a0}\ , \qquad \{E^2,\,J_{a0}\}=2EJ_{a0}\ , \tag{25}$$

$$\{J_{a0'},\,J_{b0'}\}=-J_{ab}=\{J_{a0},\,J_{b0}\}\ , \qquad \{J_{a0'},\,J_{b0}\}=E\delta_{ab}\ , \tag{26}$$

where J_{ab} have the standard form of the rotation generators

$$J_{ab}=p_a\,x^b-p_b\,x^a \tag{27}$$

and they correspond to the dynamical integrals (20) for the $\mathrm{SO}(N)$ rotations in AdS_{N+1}. Equations (25) and (26) are then equivalent to a part of the commutation relations of the symmetry group Lie algebra. The rest part of the algebra is trivially fulfilled, since the rotation generators both in AdS_{N+1} and S^M commute with the energy square (21). Thus, the Poisson bracket algebra of the dynamical integrals (20) is not deformed by the Hamiltonian reduction, which is a consequence of their gauge invariance for the initial system (1).

Concluding this section we comment on the Casimir numbers of the isometry groups. From (22) and (23) follows (see (A.4) in Appendix) that

$$\frac{1}{2}J_{AB}J^{AB}=-R^2\left(g^{00}E^2+g^{ab}p_ap_b\right), \qquad \frac{1}{2}L_{IJ}L_{IJ}=R_S^2\;g^{\alpha\beta}\pi_\alpha\pi_\beta \tag{28}$$

and due to the mass-shell condition (2) we find

$$\frac{1}{2}\,J_{AB}J^{AB}-\kappa^2\,\frac{1}{2}\,L_{IJ}\,L_{IJ}=\mathcal{M}^2\,R^2\ . \tag{29}$$

Notice that this constant coincides with E^2_{min}.

---

[2] In fact, E^2 corresponds to the energy square in units of $1/R^2$.

Quantization

In this section we first investigate the eigenvalue problem for the energy square operator. Then we construct the isometry group generators and check the algebra of their commutators. Finally, we derive the quantum version of the relation between the Casimir numbers (29).

The energy square operator is defined by (9) and in our case its eigenfunctions can be written in the following factorized form

$$\Psi = \psi(x)\, Y_L^{l_1,\cdots}(\phi)\ , \tag{30}$$

where $Y_L^{l_1,\cdots}(\phi)$ are the spherical harmonics on S^M. These states are the eigenfunctions of the Laplace-Beltrami operator Δ_S with eigenvalues $-L(L+M-1)$ for integer L.

The operator E^2 on the AdS part then is given by

$$E^2 = -h^{ab}\,\partial^2_{ab} + (N+M)x^a\,\partial_a + \frac{C}{\rho^2} + C_1\ , \tag{31}$$

with constant parameters

$$C_1 = \frac{(N+M-1)^2}{4} \tag{32}$$

and $C = \mathcal{M}^2R^2 + C_2 + C_L$, where

$$C_2 = (\kappa^2-1)\,\frac{(N+M-1)M(M-1)}{4(M+N)} \tag{33}$$

corresponds to the scalar curvature term[3] obtained by (13) and

$$C_L = \kappa^2\, L(L+M-1) \tag{34}$$

is the angular momentum contribution from rotations on S^M. The operator (31) has the same structure as in the AdS space, but with deformed parameters, which depend on the spherical part as well. Note that the pure AdS case corresponds to $M = 0 = L$.

Introducing the radial variable on the unit disk $r^2 := x^ax^a = 1-\rho^2$, we find $x^a\,\partial_a = r\,\partial_r$ and the second order derivative operator in (31) takes the form

$$h^{ab}\,\partial^2_{ab} = \partial^2_{aa} - r^2\partial^2_{rr}\ , \tag{35}$$

which is obviously invariant under the $SO(N)$ rotations in AdS_{N+1}.

Taking into account this symmetry in the operator (31), one can further factorize the eigenfunctions (30)

$$\psi(x) = F(r)\, Y_l^{l_1,\cdots}(\varphi)\ . \tag{36}$$

Here $Y_l^{l_1,\cdots}(\varphi)$ are again the spherical harmonics, but now on the unit sphere S^{N-1} $(N>1)$,[4] which can be treated as the boundary of the unit disk. Using the parametrization of the radial variable $r = \cos\sigma$, with $\sigma \in (0, \pi/2]$, and the rescaling of the radial wave function

$$F(r) = (\sin\sigma)^{-\frac{M}{2}}\,(\cos\sigma)^{-\frac{N-1}{2}}\, f(\sigma)\ , \tag{37}$$

---

[3] This term vanishes at equal radii $R = R_S$ $(\kappa = 1)$.

[4] The case $N = 1$ will be treated separately.

we find that $f(\sigma)$ satisfies the Schrödinger equation with the Pöschl-Teller potential [9]

$$\left(-\partial^2_{\sigma\sigma}+\frac{A}{\sin^2\sigma}+\frac{B}{\cos^2\sigma}\right)f(\sigma)=E^2\,f(\sigma)\ , \tag{38}$$

where

$$A=C+\frac{1}{4}\,M(M-2)\ ,\qquad B=l(l+N-2)+\frac{1}{4}\,(N-1)(N-3)\ . \tag{39}$$

The integration measure for the scalar product of wave functions $f(\sigma)$ is just $d\sigma$ and the Schrödinger operator in (38), therefore, is Hermitian.

Notice that $B\geq 3/4$, except for the two cases

$$l=0,\quad N=3,\quad B=0\ ;\qquad l=0,\quad N=2,\quad B=-1/4\ . \tag{40}$$

To have the Schrödinger operator in (38) bounded from below (see [10]), we assume $A\geq -1/4$. This condition bounds the parameter $\mathcal{M}^2R^2$.

We are looking for normalizable solutions of (38). To analyze them, it is convenient to introduce the parameters

$$\mu:=\frac{1}{2}+\sqrt{A+\frac{1}{4}}\ ,\qquad \nu:=l+\frac{N-1}{2}\ , \tag{41}$$

which are the large roots of the equations $\mu(\mu-1)=A$ and $\nu(\nu-1)=B$, respectively.

The behavior of a solution of (38) at the boundaries is given by

$$f(\sigma)\simeq c_1\,\sigma^{\mu}+c_2\,\sigma^{1-\mu}\ ,\ \ \sigma\to 0;\qquad f(\sigma)\simeq d_1\,(\pi/2-\sigma)^{\nu}+d_2\,(\pi/2-\sigma)^{1-\nu}\ ,\ \ \sigma\to\pi/2\ . \tag{42}$$

The normalizability of $f(\sigma)$ then requires that $c_2=0$ if $\mu\geq 3/2$, and $d_2=0$ if $\nu\geq 3/2$.

The case $B\geq 3/4$ is equivalent to $\nu\geq 3/2$ and the corresponding solution of (38) for $d_1=1$ and $d_2=0$ is given by

$$f(\sigma)=(\sin\sigma)^{\mu}\,(\cos\sigma)^{\nu}\ {}_2F_1\left(a,b,c;\cos^2\sigma\right)\ , \tag{43}$$

with the following parameters of the hypergeometric function

$$a=\frac{1}{2}\,(\mu+\nu-E)\ ,\qquad b=\frac{1}{2}\,(\mu+\nu+E)\ ,\qquad c=\nu+\frac{1}{2}\ . \tag{44}$$

Since the wave function (37) has to be regular at $\sigma=\pi/2$, the solution (43) describes the two exceptional cases (40) as well. To analyze the behavior of this solution at the boundary $\sigma=0$, let us consider the following two solutions of (38)

$$f_1(\sigma)=(\sin\sigma)^{\mu}\,(\cos\sigma)^{\nu}\ {}_2F_1\left(a,b,a+b-c+1;\sin^2\sigma\right)\ , \tag{45}$$

$$f_2(\sigma)=(\sin\sigma)^{1-\mu}\,(\cos\sigma)^{\nu}\ {}_2F_1\left(c-a,c-b,c-a-b+1;\sin^2\sigma\right)\ , \tag{46}$$

where $f_1(\sigma)$ corresponds to $c_1=1,\ c_2=0$ in (42) and $f_2(\sigma)$ to $c_1=0,\ c_2=1$.

The properties of hypergeometric functions then provide

$$f(\sigma) = \frac{\Gamma(c-a-b)\Gamma(c)}{\Gamma(c-a)\Gamma(c-b)}\, f_1(\sigma) + \frac{\Gamma(a+b-c)\Gamma(c)}{\Gamma(a)\Gamma(b)} f_2(\sigma)\ . \tag{47}$$

If $\mu \geq 3/2$, the normalizability condition $c_2 = 0$ requires $a = -n$ with integer n, and this condition leads to the energy spectrum

$$E_{n,l,L} = \frac{N}{2} + \sqrt{A + \frac{1}{4}} + l + 2n\ . \tag{48}$$

The dependence on L (and on the mass parameter and the radii) is contained in the parameter A (see (39) and (34)).

Let us now take $\frac{1}{2} \leq \mu < \frac{3}{2}$, which corresponds to $-\frac{1}{4} \leq A < \frac{3}{4}$. In this case the solution (43) is normalizable for any (even complex) values of E^2, which means that the Schrödinger operator in (38) is not essentially self-adjoint. However the analysis of the deficiency indices [10] shows that this operator has self-adjoint extensions. There are two different acceptable self-adjoint extensions with wave function $f_1(\sigma)$ and $f_2(\sigma)$, which are specified by the following boundary behavior at $\sigma \to 0$

$$f_1(\sigma) \simeq \sigma^{\mu}\ , \qquad f_2(\sigma) \simeq \sigma^{1-\mu}\ . \tag{49}$$

The energy spectrum in the first case is obviously given again by (48) and in the second case one finds $c - b = -n$, which is equivalent to

$$E^{(-)}_{n,l,L} = \frac{N}{2} - \sqrt{A + \frac{1}{4}} + l + 2n\ . \tag{50}$$

For $N = 1$ we use the parametrization $x^1 = -\cos\sigma$, with $\sigma \in (0, \pi)$, and the same rescaling as in (37). This leads again to the Schrödinger equation (38) with vanishing B

$$\left(-\partial^2_{\sigma\sigma} + \frac{A}{\sin^2\sigma}\right) f(\sigma) = E^2\, f(\sigma)\ . \tag{51}$$

If $A \geq 3/4$, the Schrödinger operator in (51) is essentially self-adjoint with spectrum

$$E_n^2 = (\mu + n)^2\ , \tag{52}$$

where n is non-negative integer and μ is given by (41).

If $-1/4 \leq A < 3/4$ the Schrödinger operator in (51) is not self-adjoint and one can consider two different self-adjoint extensions similarly to the higher dimensional cases.

An interesting case here is $A = 0$ (i.e. $\mu = 1$), which corresponds to a free particle in a box. The operator $-\partial^2_{\sigma\sigma}$ is then characterized by two different self-adjoint extensions. The corresponding eigenstates are give by the trigonometric functions

$$f_{1,n}(\sigma) = \sin(n+1)\sigma\ , \qquad f_{2,n}(\sigma) = \cos n\sigma\ , \quad (n \geq 0), \tag{53}$$

where $f_{1,n}$ and $f_{2,n}$ satisfy Dirichlet and Neumann boundary conditions, respectively.

Now we discuss the isometry group generators. The quantization of the rotation generators both on S^M and AdS_{N+1} is trivial. Up to the factor $(-i)$, these operators are given by the corresponding vector fields obtained from (22)-(23). Therefore, it suffices to discuss the construction of the boost generators only.

We introduce the boost operators, which correspond to the functions (24), similarly to the AdS case [1]

$$J_{a0'} = \sqrt{E}\; x^a \sqrt{E}\;, \qquad J_{a0} = i\sqrt{E}\; \left(V_a - \frac{N+M-1}{2}\, x^a\right)\, \frac{1}{\sqrt{E}}\;, \tag{54}$$

with

$$V_a := h^{ab}\partial_b = \partial_a - x^a x^b \partial_b\;. \tag{55}$$

The calculation of the commutation relations of the operators x^a and V_a with the energy square operator (31) is straightforward (see (A.7) in Appendix) and we obtain the quantum version of the Poisson bracket relations (25)

$$[E^2,\, J_{a0'}] = 2iJ_{a0}\, E + J_{a0'}\;, \qquad [E^2,\, J_{a0}] = -2iJ_{a0'}\, E + J_{a0}\;. \tag{56}$$

Note that the value (32) for the constant C_1 is important to verify the second relation in (56). This confirms the value of the coefficient of the scalar curvature term in (9).

From the commutation relations (56) follow the commutators

$$[E, Z_a] = -Z_a\;, \qquad [E, Z^*_a] = Z^*_a\;, \tag{57}$$

where $Z_a = J_{a0'} - iJ_{a0}$ and $Z^*_a = J_{a0'} + iJ_{a0}$ are the lowering and raising operators in the $\mathrm{so}(2,N)$ algebra. Then it remains to calculate the commutators between the boosts only.

In particular, by (54) one gets

$$[J_{a0'}, J_{b0'}] = \sqrt{E}\big(x^a\, E\, x^b\, E - x^b\, E\, x^a\, E\big)\frac{1}{\sqrt{E}}\;. \tag{58}$$

To simplify the right hand side here, we apply the same trick as in [1]. Using the commutation relation $[E, J_{a0'}] = iJ_{a0}$, which follows from (57), one finds the operator equality

$$E\, x^a\, E - x^a\, E^2 = \frac{N+M-1}{2}\, x^a - V_a\;. \tag{59}$$

The operator in the parentheses of (58), therefore, corresponds to the generator of $\mathrm{SO}(N)$ rotations (27) given by $iJ_{ab} = x^b\partial_a - x^a\partial_b$. Since $[E, J_{ab}] = 0$, one can neglect the $\sqrt{E}$ terms in (58) and obtain

$$[J_{a0'}\,,\, J_{b0'}] = iJ_{ab}. \tag{60}$$

The other commutators between the boost operators are computed in a similar way and they realize the algebra (26) on the quantum level.

The lowering and raising operators Z_a and Z^*_a provide an alternative way for calculation of the energy spectrum. One can start with the ground state ψ_0 of the operator (31). This state is a $\mathrm{SO}(N)$ scalar and it is annihilated by the lowering operators

$$Z_a = \sqrt{E}\; \left(V_a + x^a\Big(E - \frac{N+M-1}{2}\Big)\right)\, \frac{1}{\sqrt{E}}\;. \tag{61}$$

The ground state wave function $\psi_0(\rho)$ then satisfies the equations

$$V_a\,\psi_0(\rho) = x^a \left(\frac{N+M-1}{2} - E_0\right)\psi_0(\rho)\ , \quad a = 1, ...N, \tag{62}$$

where E_0 denotes the energy of the ground state. Using (A.6), we find the solution

$$\psi_0(\rho) = c_0\,\rho^{E_0 - \frac{N+M-1}{2}}\ , \tag{63}$$

up to a normalization constant c_0. The normalizability condition with integration measure (16) restricts the minimal energy by the unitarity bound in the AdS space [11]

$$E_0 > \frac{N}{2} - 1\ . \tag{64}$$

The wave function (63) should also be an eigenfunction of the energy square operator (31) with eigenvalue E_0^2. This condition relates E_0 to other parameters of the theory in the form

$$\left(E_0 - \frac{N}{2}\right)^2 = A + \frac{1}{4}\ , \tag{65}$$

where A is given by (39). If $1/4 \leq A < 3/4$ one gets two solutions of (65) as above. However, since A is unbounded for increasing L, one has to take only the large root E_0^+ of (65). The action of the raising operators on the ground state shifts the energy levels by one and we obtain the same spectrum as above and the following representation of the isometry group

$$\sum_{L\geq 0} D_{E_0^+(L)}(\mathrm{O}(2,N)) \otimes D_L(\mathrm{O}(M+1))\ , \tag{66}$$

where $D_{E_0^+(L)}(\mathrm{O}(2,N))$ and $D_L(\mathrm{O}(M+1))$ are the corresponding standard representations.

Taking the case of equal radii ($\kappa = 1$) and $M = N+1$, we obtain

$$E_0^+(L) = \frac{N}{2} + \sqrt{\mathcal{M}^2 R^2 + L(L+N) + \frac{N^2}{4}}\ . \tag{67}$$

Finally, we consider the quantum version of the relation (29) between the Casimir numbers. In the calculation of the Casimir number operator of the SO$(2,N)$ group one can use the identity (59) for the term $-J_{a0'}\,J_{a0'}$ and the factors $\sqrt{E}$ in the term $-(J_{a0'}\,J_{a0'} + J_{a0}\,J_{a0})$ can be removed by the same trick as in (58). The computation of the remaining part is straightforward, and taking into account that $\frac{1}{2}\,L_{IJ}\,L_{IJ} = -\Delta_S$, we find

$$\frac{1}{2}\,J_{AB}\,J^{AB} - \kappa^2\,\frac{1}{2}\,L_{IJ}\,L_{IJ} = \mathcal{M}^2\,R^2 + C_2 - \frac{N^2-(M-1)^2}{4}\ . \tag{68}$$

Notice that the quantum correction here vanishes for $\kappa = 1$ (i.e. $R = R_s$) and $N+1 = M$.

Acknowledgments

We thank Harald Dorn for his interest in our work and for useful comments. G.J. thanks Martin Reuter for helpful discussions on static gauge quantization. This work was done during the visit of G.J. and Z.K. at Humboldt University of Berlin. They thank the department of Physics for warm hospitality.
This work has been supported in part by the grant I/84600 from VolkswagenStiftung.

Appendix

In this Appendix we collect some useful formulae used in the main text.

Weyl transformation of a scalar curvature

If a metric tensor $h_{mn}(x)$ is a Weyl transformed $g_{mn}(x)$

$$h_{mn}(x) = \frac{1}{\Lambda(x)}\, g_{mn}(x) \ ,$$

then its scalar curvature is given by

$$\mathcal{R}_h = \Lambda\left(\mathcal{R}_g + (\mathcal{N}-1)\Delta_g(\log\Lambda) - \frac{(\mathcal{N}-1)(\mathcal{N}-2)}{4}\, g^{mn}\,\partial_m(\log\Lambda)\,\partial_n(\log\Lambda)\right) \ . \qquad \text{(A.1)}$$

Killing vector fields on AdS$_{N+1}$ and S^M

The Killing vector fields on AdS$_{N+1}$ and S^M are treated similarly and it is sufficient to consider the spherical part only.

Let us introduce vector fields $\mathcal{V}_I$ on S^M with components $\mathcal{V}_I^\alpha = g^{\alpha\beta}\partial_\beta Y_I$. These fields are obtained from the equations $\mathcal{V}_I \rfloor g = \mathrm{dY}_I$ and they satisfy the relations

$$\mathcal{V}_I(Y_K) = \delta_{IK} - \frac{Y_I\, Y_K}{R^2} \ . \qquad \text{(A.2)}$$

The Killing vector fields $\mathcal{V}_{IJ}$ with components (23) then can be written as $\mathcal{V}_{IJ} = Y_J\,\mathcal{V}_I - Y_I\,\mathcal{V}_J$, and due to (A.2) they provide the infinitesimal form of the isometry transformations

$$\mathcal{V}_{IJ}(Y_K) = \delta_{IK}\, Y_J - \delta_{JK}\, Y_I \ . \qquad \text{(A.3)}$$

The metric on S^M can be written as $\mathrm{dY}_I\,\mathrm{dY}_I = \frac{1}{2}\,\theta_{IJ}\,\theta_{IJ}$, with $\theta_{IJ} = Y_I\,\mathrm{dY}_I - Y_I\,\mathrm{dY}_J$, and in local coordinates one finds

$$\frac{1}{2}\,\mathcal{V}_{IJ}^\alpha\,\mathcal{V}_{IJ}^\beta = g^{\alpha\beta} \ . \qquad \text{(A.4)}$$

Algebra of the boost operators

The energy square operator (31) is linear in $H = -h^{ab}\partial^2_{ab}$ and $D = x^a\partial_a$, whereas the boost generators (54) are linear in V_a and in the multiplication operator x^a. The commutators of these operators form the following algebra

$$\begin{aligned} &[D, x^a] = x^a, && [H, x^a] = -2V_a \\ &[D, V_a] = -2x^a D - V_a \ , \qquad && [H, V_a] = 2V_a - 2x^a H \ , \end{aligned} \qquad \text{(A.5)}$$

and, in addition, one has

$$[V_a, \rho] = V_a(\rho) = -x^a \rho \ . \qquad \text{(A.6)}$$

From these commutators follow

$$[E^2, x^a] = -2V_a + (M+N)x^a \ , \qquad [E^2, V_a] = -2x^a(E^2 - C_1) - (M+N-2)V_a \ , \qquad \text{(A.7)}$$

which turn the equations in (56) to identities.



Figure 1: *The unit disk here is tangent to the Lobachevski plane at the pole. The coordinates on the disk used in the parametrization* (10) *correspond to the stereographic projection onto the disk made from the origin of the 3d space. The unit half-sphere is also tangent to the disk at the pole. The same coordinates parameterize the unit half-sphere by the vertical projection.*

References

[1] H. Dorn, G. Jorjadze, C. Kalousios and J. Plefka, *"Coordinate representation of particle dynamics in AdS and in generic static spacetimes,"* J. Phys. A A **44** (2011) 095402 [arXiv:1011.3416].

[2] L. D. Faddeev and R. Jackiw, *"Hamiltonian Reduction of Unconstrained and Constrained Systems,"* Phys. Rev. Lett. **60** (1988) 1692.

[3] G. Jorjadze, J. Plefka and J. Pollok, *"Bosonic String Quantization in Static Gauge,"* [arXiv:1207.4368].

[4] B. S. DeWitt, *"Point Transformations In Quantum Mechanics,"* Phys. Rev. **85**, 653 (1952). • B. S. DeWitt, *"Dynamical theory in curved spaces. 1. A Review of the classical and quantum action principles,"* Rev. Mod. Phys. **29**, 377 (1957).

[5] F. Bastianelli and P. van Nieuwenhuizen, *"Path integrals and anomalies in curved space,"* Cambridge, UK: Univ. Pr. (2006) 379 P

[6] O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri and Y. Oz, *"Large N field theories, string theory and gravity,"* Phys. Rept. **323**, 183 (2000) [arXiv:hep-th/9905111].

[7] H. Dorn, M. Salizzoni and C. Sieg, *"On the propagator of a scalar field on AdS×S and on the BMN plane wave,"* JHEP **0502** (2005) 047 [hep-th/0307229].

[8] F. Passerini, J. Plefka, G. W. Semenoff and D. Young, *"On the Spectrum of the $AdS_5 \times S^5$ String at large lambda,"* JHEP **1103** (2011) 046 [arXiv:1012.4471].

[9] G. Pöschl and E. Teller, *"Bemerkungen zur Quantenmechanik des anharmonischen Oszillators,"* Z. Physik **83**, 143 (1933).

[10] M. Reed and B. Simon, *"Methods Of Modern Mathematical Physics. Vol. 2: Fourier Analysis, Self-Adjointness,"* New York, USA: Academic Press (1975), 361 P.

[11] P. Breitenlohner and D. Z. Freedman, *"Stability In Gauged Extended Supergravity,"* Annals Phys. **144**, 249 (1982).